

**AMENDMENTS TO THE CLAIMS**

1. (Currently amended) A method of treating neuromuscular dysfunction of the lower urinary tract in a mammal in need of such treatment comprising administering to said mammal an effective amount of a compound having selective affinity for the mGlu5 subtype of the metabotropic glutamate receptors to treat said neuromuscular dysfunction of the lower urinary tract.

2. (Previously presented) The method of claim 1 wherein said compound has an at least about 10-fold selectivity for the mGlu5 subtype of the metabotropic glutamate receptors.

3. (Currently amended) The method of claim 1 wherein said compound has an at least about 25-fold ~~slectivity~~ selectivity for the mGlu5 subtype of the metabotropic glutamate receptors.

4. (Currently amended) The method of claim 1 wherein said compound has an at least about 50-fold ~~slectivity~~ selectivity for the mGlu5 subtype of the metabotropic glutamate receptors.

5. (Currently amended) The method of claim 1 wherein said compound has an at least about 100-fold ~~slectivity~~ selectivity for the mGlu5 subtype of the metabotropic glutamate receptors.

6. (Currently amended) The method of claim 1 wherein said compound has an at least about 500-fold ~~slectivity~~ selectivity for the mGlu5 subtype of the metabotropic glutamate receptors.

7. (Original) The method of claim 1 wherein said compound is a selective mGlu5 receptor antagonist.

8. (Original) The method of claim 7 wherein said neuromuscular dysfunction is urinary urgency, overactive bladder, increased urinary frequency, decreased urinary compliance, cystitis, incontinence, urine leakage, enuresis, dysuria, urinary hesitancy or difficulty in emptying the bladder.

9. (Withdrawn) The method of claim 8 wherein said neuromuscular dysfunction that is decreased urinary compliance is decreased bladder storage capacity.

10. (Withdrawn) The method of claim 8 wherein said neuromuscular dysfunction is interstitial cystitis.

11. (Original) The method of claim 1 wherein said compound is administered as a pharmaceutically acceptable composition.

12. (Original) The method of claim 11 wherein said compound is administered via an oral, parenteral, intranasal, sublingual, rectal or inhalatory route, or by insufflation, transdermal patches or lyophilized composition.

13. (Original) The method of claims 1 wherein said compound is administered in an amount of between about 0.01 to about 25 mg/kg/day.

14. (Original) The method of claim 13 wherein said compound is administered in an amount of between about 0.1 to about 10 mg/kg/day.

15. (Original) The method of claim 14 wherein said compound is administered in an amount of about 0.2 to about 5 mg/kg/day.

16. (Original) The method of claim 1 wherein said compound is administered at a total daily dose of about 25 to about 1000 mg.

17. (Original) The method of claim 16 wherein said compound is administered at a total daily dose of about 150 to about 500 mg.

18. (Original) The method of claim 17 wherein said compound is administered at a total daily dose of about 350 mg.

19. (Original) The method of claim 1 wherein said compound is administered in combination with an antimuscarinic drug.

20. (Original) The method of claim 19 wherein said antimuscarinic drug is selected from the group consisting of oxybutynin, tolterodine, darifenacin and temiverine.

21. (Withdrawn) The method of claim 1 wherein said compound is administered in combination with an  $\alpha$ 1-adrenergic antagonist.

22. (Withdrawn) The method of claim 21 wherein said  $\alpha$ 1-adrenergic antagonist is selected from the group consisting of prazosin, doxazosin, terazosin, alfuzosin and tamsulosin.

23. (Withdrawn) The method of claim 1 wherein said compound is administered in combination with a 5-HT<sub>1A</sub> receptor antagonist.

24. (Withdrawn) The method of claim 1 wherein said compound is administered in combination with a selective COX2 inhibitor.

25. (Withdrawn) The method of claim 24 wherein said selective COX2 inhibitor comprises a NO releasing group.

26. (Withdrawn) The method of claim 1 wherein said compound is administered in combination with a non-selective COX1/COX2 inhibitor.


27. (Withdrawn) The method of claim 26 wherein said non-selective COX1/COX2 inhibitor derivative comprises a NO releasing group.

28. (Original) The method of claim 1 wherein said mammal is a human.

29. (Original) The method of claim 1 wherein said compound is administered in admixture with a pharmaceutically acceptable diluent or carrier.

30. (Original) The method of claim 29 wherein said pharmaceutically acceptable diluent or carrier is selected from the group consisting of ethanol, water, glycerol, aloe vera gel, allantoin, glycerine, vitamin A oil, vitamin E oil, mineral oil, phosphate buffered saline, PPG2 myristyl propionate, magnesium carbonate, potassium phosphate, vegetable oil, animal oil, and solketal.

31. (Withdrawn) The method of claim 1 wherein said compound having selective affinity for the mGlu5 subtype of the metabotropic glutamate receptors has a general formula I



wherein:

R<sub>1</sub> represents hydrogen, lower alkyl, lower hydroxyalkyl, lower alkylamino, piperidino, carboxyl, esterified carboxyl, amidated carboxyl, lower alkoxy, lower haloalkyl, lower haloalkoxy, cyano, alkynyl, lower alkoxycarbonyl, di-(lower)alkylamino, lower alkylaminocarbonyl, trifluoromethylphenylaminocarbonyl or N-(lower)alkyl-N-phenylcarbamoyl, said N-(lower)alkyl and N-phenyl radicals being unsubstituted or substituted independently with a substituent selected from the group consisting of lower alkyl, lower alkoxy, halogen, and trifluoromethyl groups,

R<sub>2</sub> represents hydrogen, lower alkyl, carboxyl, esterified carboxyl, amidated carboxyl, lower hydroxyalkyl, hydroxyl, lower alkoxy or lower alkanoyloxy, lower alkoxycarbonyl, di-(lower)-alkylamino-(lower)alkanoyl, di-(lower)alkylaminomethyl, 4-(4-fluorobenzoyl)-piperidin-1-yl-




or an enantiomer, diastereoisomer, N-oxide, crystalline form, hydrate, solvate, pharmacologically active metabolite, prodrug, or pharmaceutically acceptable salt thereof.

R<sub>1</sub> is hydrogen, (C<sub>1-4</sub>)alkyl, (C<sub>1-4</sub>)alkoxy, hydroxy(C<sub>1-4</sub>)alkyl, cyano, ethynyl, carboxy, (C<sub>1-4</sub>)alkoxycarbonyl, di(C<sub>1-4</sub>)alkylamino, (C<sub>1-6</sub>)alkylaminocarbonyl, or trifluoromethylphenylaminocarbonyl;

R<sub>2</sub> is hydrogen, hydroxy, (C<sub>1-4</sub>) alkyl, hydroxy (C<sub>1-4</sub>) alkyl, (C<sub>1-4</sub>) alkoxy, carboxy, (C<sub>2-5</sub>)alkanoyloxy, (C<sub>1-4</sub>)alkoxycarbonyl, di(C<sub>1-4</sub>)alkylamino(C<sub>1-4</sub>)alkanoyl, di(C<sub>1-4</sub>)alkylaminomethyl, 4-(4-fluorobenzoyl)-piperidin-1-yl-carbonyl, 4-*tert*-butyloxycarbonyl-piperazin-1-yl-carbonyl, 4-(4-azido-2-hydroxybenzoyl)-piperazin-1-yl-carbonyl or 4-(4-azido-2-hydroxy-3-iodobenzoyl)-piperazin-1-yl-carbonyl;

R<sub>4</sub> is hydrogen, hydroxy, (C<sub>1-4</sub>)alkoxy, carboxy, (C<sub>2-5</sub>)alkanoyloxy, (C<sub>1-4</sub>)alkoxycarbonyl, amino(C<sub>1-4</sub>)alkoxy, di(C<sub>1-4</sub>)alkylamino(C<sub>1-4</sub>)alkoxy, di(C<sub>1-4</sub>)alkylamino(C<sub>1-4</sub>)alkyl, carboxy(C<sub>1-4</sub>)alkylcarbonyl, (C<sub>1-4</sub>)alkoxycarbonyl(C<sub>1-4</sub>)alkoxy, hydroxy(C<sub>1-4</sub>)alkyl, di(C<sub>1-4</sub>)alkylamino(C<sub>1-4</sub>)alkoxy, or m-hydroxy-p-azidophenylcarbonylamino (C<sub>1-4</sub>)alkoxy; and

$R_5$  is a group of formula



R<sub>a</sub> and R<sub>b</sub> independently are hydrogen, hydroxy, halogen, nitro, cyano, carboxy, (C<sub>1-4</sub>)alkyl, (C<sub>1-4</sub>)alkoxy, hydroxy(C<sub>1-4</sub>)alkyl, (C<sub>1-4</sub>)alkoxycarbonyl, (C<sub>2-7</sub>)alkanoyl, (C<sub>2-5</sub>)alkanoyloxy, (C<sub>2-5</sub>)alkanoyloxy(C<sub>1-4</sub>)alkyl, trifluoromethyl, trifluoromethoxy, trimethylsilylethynyl, (C<sub>2-5</sub>)alkynyl, amino, azido, amino(C<sub>1-4</sub>)alkoxy, (C<sub>2-5</sub>)alkanoylamino(C<sub>1-4</sub>)alkoxy, (C<sub>1-4</sub>)alkylamino(C<sub>1-4</sub>)alkoxy, di(C<sub>1-4</sub>)alkylamino(C<sub>1-4</sub>)alkoxy, (C<sub>1-4</sub>)alkylamino, di(C<sub>1-4</sub>)alkylamino, monohalobenzylamino, thienylmethylamino, thienylcarbonylamino, trifluoromethylphenylaminocarbonyl, tetrazolyl, (C<sub>2-5</sub>)alkanoylamino, benzylcarbonylamino, (C<sub>1-4</sub>)alkylaminocarbonylamino (C<sub>1-4</sub>)alkoxycarbonylaminocarbonylamino or (C<sub>1-4</sub>)alkylsulfonyl;

R<sub>c</sub> is hydrogen, fluorine, chlorine, bromine, hydroxy, (C<sub>1-4</sub>)alkyl, (C<sub>2-5</sub>)alkanoyloxy, (C<sub>1-4</sub>)alkoxy or cyano; and

R<sub>d</sub> is hydrogen, halogen or (C<sub>1-4</sub>)alkyl.

33. (Withdrawn) The method of claim 31 wherein said compound has a structure wherein X<sub>1</sub> is a (C<sub>2-4</sub>)alkenylene, (C<sub>2-4</sub>)haloalkenylene, (C<sub>2-4</sub>)alkynylene or (C<sub>2-4</sub>)haloalkynylene group, wherein each of the foregoing groups is linked via vicinal unsaturated carbon atoms;


R<sub>1</sub> is hydrogen, (C<sub>1-4</sub>)alkyl, (C<sub>1-4</sub>)alkoxy, cyano, ethynyl or di(C<sub>1-4</sub>)alkylamino;

R<sub>2</sub> is hydrogen, hydroxy, carboxy, (C<sub>1-4</sub>)alkoxycarbonyl, di(C<sub>1-4</sub>)alkylaminomethyl, 4-(4-fluorobenzoyl)-piperidin-1-yl-carbonyl, 4-*tert*-butyloxycarbonyl-piperazin-1-yl-carbonyl, 4-(4-azido-2-hydroxybenzoyl)-piperazin-1-yl-carbonyl or 4-(4-azido-2-hydroxy-3-iodobenzoyl)-piperazin-1-yl-carbonyl;

R<sub>3</sub> is hydrogen, (C<sub>1-4</sub>)alkyl, carboxy, (C<sub>1-4</sub>)alkoxycarbonyl, (C<sub>1-4</sub>)alkylcarbamoyl, hydroxy(C<sub>1-4</sub>)alkyl, di(C<sub>1-4</sub>)alkylaminomethyl, morpholinocarbonyl or 4-(4-fluoro-benzoyl)-piperidin-1-yl-carbonyl;

R<sub>4</sub> is hydrogen, hydroxy, carboxy, (C<sub>2-5</sub>)alkanoyloxy, (C<sub>1-4</sub>)alkoxycarbonyl, amino(C<sub>1-4</sub>)alkoxy, di(C<sub>1-4</sub>)alkylamino(C<sub>1-4</sub>)alkoxy, di(C<sub>1-4</sub>)alkylamino(C<sub>1-4</sub>)alkyl or hydroxy(C<sub>1-4</sub>)alkyl; and

R<sub>5</sub> is a group of formula



R<sub>a</sub> and R<sub>b</sub> independently are hydrogen, halogen, nitro, cyano, (C<sub>1-4</sub>)alkyl, (C<sub>1-4</sub>)alkoxy, trifluoromethyl, trifluoromethoxy or (C<sub>2-5</sub>)alkynyl;

R<sub>c</sub> is hydrogen, fluorine, chlorine, bromine, hydroxy, (C<sub>1-4</sub>)alkyl, (C<sub>2-5</sub>)alkanoyloxy, (C<sub>1-4</sub>)alkoxy or cyano; and


R<sub>d</sub> is hydrogen, halogen or (C<sub>1-4</sub>)alkyl.



34. (Withdrawn) The method of claim 31 wherein said compound is 2-methyl-6-(phenylethynyl)pyridine (MPEP).

35. (Withdrawn) The method of claim 31 wherein said compound is 2-methyl-6-(2-phenylethenyl)pyridine (SIB 1893).


36. (Withdrawn) The method of claim 1 wherein said compound has a general formula I-A



wherein

R' is hydrogen or (C<sub>1-4</sub>)alkyl and

M is a group of formula



wherein

R<sub>aa</sub>, R<sub>bb</sub> and R<sub>cc</sub> are independently of each other hydrogen, (C<sub>1-4</sub>)alkyl, (C<sub>1-4</sub>)alkoxy, hydroxyl, (C<sub>1-4</sub>)hydroxyalkyl, cyano or halo,

R<sub>dd</sub> is cyano or halo,

R<sub>ee</sub> is hydroxyl, (C<sub>1-4</sub>)alkyl or (C<sub>1-4</sub>)alkoxy,

R<sub>ff</sub> is hydrogen or (C<sub>1-4</sub>)alkyl,

R<sub>ij</sub> and R<sub>ji</sub> are independently hydrogen, (C<sub>1-4</sub>)alkyl or phenyl, and


or an enantiomer, diastereoisomer, N-oxide, crystalline form, hydrate, solvate,

R7C1C(R6)C(B1#C)C(R10)C(R9)C(R8)C1

II-A

R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub> and R<sub>10</sub> represent, independently from each other, hydrogen, lower alkyl, lower alkoxy, -(CH<sub>2</sub>)<sub>n</sub>-halo, -(CH<sub>2</sub>)<sub>n</sub>-NR<sub>e</sub>R<sub>f</sub>, -(CH<sub>2</sub>)<sub>n</sub>-N(R<sub>e</sub>)-C(O)-(lower)alkyl, aryl or heteroaryl, which is unsubstituted or substituted by one or more lower alkyl groups;

B<sub>1</sub> represents



wherein

- R<sub>11</sub> represents hydrogen, lower alkyl, -(CH<sub>2</sub>)<sub>n</sub>-C(O)OR<sub>e</sub> or halo;
- R<sub>12</sub> represents hydrogen, lower alkyl, -(CH<sub>2</sub>)<sub>n</sub>-C(O)OR<sub>f</sub>, halo, nitro or heteroaryl which is unsubstituted or substituted with lower alkyl or cycloalkyl;
- R<sub>13</sub> represents hydrogen, lower alkyl, -(CH<sub>2</sub>)<sub>n</sub>-OH, -(CH<sub>2</sub>)<sub>n</sub>-C(O)OR<sub>g</sub> or aryl;
- R<sub>14</sub> represents lower alkyl;
- R<sub>15</sub> represents hydrogen, lower alkyl or halo;
- R<sub>16</sub> represents hydrogen or alkyl;
- R<sub>17</sub> represents -(CH<sub>2</sub>)<sub>n</sub>-N(R<sub>e</sub>)-C(O)-lower alkyl;
- R<sub>18</sub> represents hydrogen or lower alkyl;
- R<sub>19</sub>, R<sub>20</sub>, R<sub>21</sub> and R<sub>22</sub> represent, independently from each other, hydrogen, lower alkyl, -(CH<sub>2</sub>)<sub>n</sub>-halo or lower alkoxy;

or an enantiomer, diastereoisomer, N-oxide, crystalline form, hydrate, solvate, pharmacologically active metabolite, prodrug, or pharmaceutically acceptable salt thereof.

40. (Withdrawn) The method of claim 1 wherein said compound has general formula II-B or II-C



R<sub>32</sub>, R<sub>33</sub>, R<sub>34</sub>, R<sub>35</sub> and R<sub>36</sub> represent, independently from each other, hydrogen, lower alkyl, -(CH<sub>2</sub>)<sub>n</sub>-halogen, lower alkoxy, -(CH<sub>2</sub>)<sub>n</sub>-NR<sub>e</sub>R<sub>f</sub>, -(CH<sub>2</sub>)<sub>n</sub>-N(R<sub>e</sub>)-C(O)-(lower)alkyl, aryl or heteroaryl which is unsubstituted or substituted by one or more lower alkyl residues;

R<sub>38</sub> represents hydrogen, lower alkyl, -(CH<sub>2</sub>)<sub>n</sub>-C(O)OR<sub>f</sub>, halogen, nitro or heteroaryl which is unsubstituted or substituted with lower alkyl or cycloalkyl;


R<sub>41</sub> represents hydrogen, halogen or lower alkyl; and

R<sub>e</sub>, R<sub>f</sub> and R<sub>g</sub> represent, independently from each other, hydrogen or lower alkyl; and

or an enantiomer, diastereoisomer, N-oxide, crystalline form, hydrate, solvate, pharmacologically active metabolite, prodrug, or pharmaceutically acceptable salt thereof.

$$A_1\text{---}L_1\text{---}B_2 \quad \text{III}$$

**A<sub>1</sub> is a 5-, 6- or 7-membered ring having the structure**



W, X<sub>3</sub>, Y<sub>2</sub> and Z<sub>1</sub> together form a 3-5 membered chain, at least one of W, X<sub>3</sub>, Y<sub>2</sub> and Z<sub>1</sub> is a group (CR<sub>h</sub>)<sub>p</sub>, wherein p is 1 or 2; and the remainder of W, X<sub>3</sub>, Y<sub>2</sub> and Z<sub>1</sub> are each independently present or absent and when present are each independently C, O, N or S;

each  $R_n$  is independently, halogen, substituted or unsubstituted hydrocarbyl, substituted or unsubstituted aryl, substituted or unsubstituted heterocyclic, substituted or unsubstituted lower alkoxy, (lower)alkylcarbonyloxy, carboxyl, esterified carboxyl, amidated carboxyl, substituted or unsubstituted lower alkylthio, substituted or unsubstituted cycloalkyl, mercapto, nitro, carboxyl, carbamate, carboxamide, hydroxyl, ester, cyano, amine, amide, amidine, amido, sulfonyl, sulfonamide or N-(lower)-alkyl-N-phenylcarbamoyl wherein each nitrogen atom is independently unsubstituted or substituted independently with lower alkyl, lower alkoxy, halo or trifluoromethyl and wherein q is 0, 1, 2 or 3;

$L_1$  is substituted or unsubstituted alkenyl, alkynyl, or azo; and


$B_2$  is substituted or unsubstituted hydrocarbyl, substituted or unsubstituted cyclohydrocarbyl, substituted or unsubstituted heterocyclic, optionally containing one or more double bonds, or substituted or unsubstituted aryl,

wherein "substituted" refers to a radical wherein one or more hydrogen atoms has been replaced with a substituent selected from the group consisting of hydroxyl, alkyl, alkoxy, mercapto, aryl, heterocycle, halogen, trifluoromethyl, pentafluoroethyl, cyano, cyanomethyl, nitro, amino, N-substituted- or N,N-di-substituted amino, wherein one or both nitrogen atoms are substituted independently with alkyl, heterocycle, aryl which are each optionally further substituted independently with hydroxyl, alkyl or heterocycle, or, alkylamide, amidine, amido, carboxy, esterified carboxy, amidated carboxy, carboxamide, carbamate, ester, sulfonyl and sulfonamide groups, and the like,

or an enantiomer, diastereoisomer, N-oxide, crystalline form, hydrate, solvate, pharmacologically active metabolite, prodrug, or pharmaceutically acceptable salt thereof.

42. (Original) The method of claim 41 wherein said administered compound is 3-(2-methylthiazol-4-yl)ethynylpyridine (MTEP).

43. (Withdrawn) The method of claim 1 wherein said compound has a general formula IV



wherein,

n is 0, 1 or 2;

X<sub>4</sub> is O, S, NH, or NOH;


R<sub>43</sub> and R<sub>44</sub> are each independently hydrogen, CN, COOR<sub>i</sub>, CONHR<sub>i</sub>, (C<sub>1-6</sub>)alkyl, or tetrazole, or R<sub>43</sub> and R<sub>44</sub> together represent an oxo group;

R<sub>i</sub> is hydrogen or (C<sub>1-6</sub>)alkyl;

R<sub>45</sub> is (C<sub>1-6</sub>)alkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-8</sub>)cycloalkyl, -CH<sub>2</sub>OH, -CH<sub>2</sub>O-alkyl, or -COOH;

Ar<sub>1</sub> is an unsubstituted aromatic or heteroaromatic group or an aromatic or heteroaromatic group substituted with one or more substituent selected from the group consisting of (C<sub>1-6</sub>)alkylamino, di-(C<sub>1-6</sub>)-alkylamino, (C<sub>1-6</sub>)alkoxy, carboxy, hydroxyl, cyano, halo, trifluoromethyl, nitro, amino, (C<sub>1-6</sub>)acylamino, (C<sub>1-6</sub>)alkylthio, (C<sub>1-6</sub>)hydroxyalkyl, (C<sub>1-6</sub>)alkylsulfonyl, and (C<sub>1-6</sub>)haloalkyl;

Z<sub>2</sub> represents a group of the formula




wherein,

R<sub>46</sub> and R<sub>47</sub> are each independently from each other hydrogen, halogen, (C<sub>1-6</sub>)alkoxy, -OAr<sub>1</sub>, (C<sub>1-6</sub>)alkyl, -CF<sub>3</sub>, COOR<sub>i</sub>, CONHR<sub>i</sub>, -CN, -OH, COR<sub>i</sub>, -S-(C<sub>1-6</sub>)-alkyl, or -SO<sub>2</sub>-(C<sub>1-6</sub>)-alkyl;

or an enantiomer, diastereoisomer, N-oxide, crystalline form, hydrate, solvate, pharmacologically active metabolite, prodrug, or pharmaceutically acceptable salt thereof.

44. (Withdrawn) The method of claim 1 wherein said compound has general formula V-A



wherein

Ar<sub>2</sub> is a heteroaryl group,

Ar<sub>3</sub> is an aryl group, where

Ar<sub>2</sub> and Ar<sub>3</sub> are each independently of each other optionally substituted with one or more substituents selected from the group consisting of -F, -Cl, -Br, -I, -OR<sub>j</sub>, -SR<sub>j</sub>, -SOR<sub>j</sub>, -SO<sub>2</sub>R<sub>j</sub>, -SO<sub>2</sub>NR<sub>j</sub>R<sub>k</sub>, -OCOR<sub>j</sub>, -OCONR<sub>j</sub>R<sub>k</sub>, -NRCOR<sub>k</sub>, -NRCO<sub>2</sub>R<sub>k</sub>, -CN, -NO<sub>2</sub>, -CO<sub>2</sub>R<sub>j</sub>, -CONR<sub>j</sub>R<sub>k</sub>, -C(O)R<sub>j</sub>, -CH(OR<sub>j</sub>)R<sub>k</sub>, -CH<sub>2</sub>(OR<sub>j</sub>), -R<sub>j</sub>, and -A-(CH<sub>2</sub>)<sub>n</sub>-NR<sub>j</sub>R<sub>k</sub>, wherein R<sub>j</sub> and R<sub>k</sub> are selected independently from the group consisting of H, CF<sub>3</sub>, (C<sub>1-10</sub>)alkyl, cycloalkyl, alkyl-aryl, alkyl-heteroaryl, heterocycloalkyl, aryl, or R<sub>j</sub> and R<sub>k</sub> may combine to form a C<sub>1-5</sub> methylene chain, and A is defined as CH<sub>2</sub>, O, NH, S, SO, SO<sub>2</sub> and n is 1, 2, 3, or 4,

G<sub>1</sub> is selected from the group consisting of -NH-, -S-, -O-, -CO-, -CONH-, -CONHCH<sub>2</sub>-, -CH<sub>2</sub>CONH-, -CH<sub>2</sub>NHNH-, -CH<sub>2</sub>NHNHCH<sub>2</sub>-, -C=NO-CH<sub>2</sub>-, -CH<sub>2</sub>NHCH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>NH-, -NHCH<sub>2</sub>CO-, -NHCH<sub>2</sub>CHOH-, -NHCH<sub>2</sub>NHNH-, -NHCONH-, or G<sub>1</sub> is a cyclic group selected from the group consisting of cyclopentane, cyclopentadiene, furan, thiofuran, pyrrolidine, pyrrole, 2-imidazoline, 3-imidazoline, 4-imidazoline, imidazole, pyrazoline, pyrazolidine, imidazolidine, oxazole, 2-oxazole, thiazole, isoxazole, isothiazole, 1*H*-1,2,4-triazole, 1*H*-1,2,3-triazole, 1,2,4-oxathiazole, 1,3,4-oxathiazole, 1,4,2-dioxazole, 1,4,2-oxathiazole, 1,2,4-oxadiazole, 1,2,4-thiadiazole, 1,2,5-oxadiazole, 1,2,5-thiadiazole, 1,3,4-oxadiazole, 1,3,4-thiadiazole, 1*H*-tetrazole, cyclohexane, piperidine, tetrahydropyridine, 1,4-dihydropyridine, pyridine, benzene,



tetrahydropyran, 3,4-dihydro-2*H*-pyran, 2*H*-pyran, 4*H*-pyran, tetrahydrothiopyran, 3,4-dihydro-2*H*-thiopyran, 2*H*-thiin, 4*H*-thiopyran, morpholine, thiomorpholine, piperazine, pyridazine, pyrimidine, pyrazine, 1,2,4-triazine, 1,2,3-triazine, 1,3,5-triazine, and 1,2,4,5-tetrazine groups,


or an enantiomer, diastereoisomer, N-oxide, crystalline form, hydrate, solvate, pharmacologically active metabolite, prodrug, or pharmaceutically acceptable salt thereof.

45. (Withdrawn) The method of claim 44 wherein Ar<sub>3</sub> is selected from the group consisting of phenyl, benzyl, naphthyl, fluorenyl, anthrenyl, indenyl, phenanthrenyl and benzonaphthenyl groups.

46. (Withdrawn) The method of claim 44 wherein Ar<sub>2</sub> is selected from the group consisting of thiazolyl, furyl, pyranyl, 2*H*-pyrrolyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, benzothiazolyl, benzimidazolyl, 3*H*-indolyl, indolyl, indazolyl, purinyl, quinoliziny, isoquinolyl, quinolyl, phthaliziny, naphthyridinyl, quinazolinyl, cinnolinyl, isothiazolyl, quinoxalinyl, indoliziny, isoindolyl, benzothienyl, benzofuranyl, isobenzofuranyl and chromenyl groups.

47. (Withdrawn) The method of claim 44 wherein Ar<sub>3</sub> is selected from the group consisting of phenyl, benzyl, naphthyl, fluorenyl, anthrenyl, indenyl, phenanthrenyl and benzonaphthenyl groups and Ar<sub>2</sub> is selected from the group consisting of thiazolyl, furyl, pyranyl, 2*H*-pyrrolyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, benzothiazolyl, benzimidazolyl, 3*H*-indolyl, indolyl, indazolyl, purinyl, quinoliziny, isoquinolyl, quinolyl, phthaliziny, naphthyridinyl, quinazolinyl, cinnolinyl, isothiazolyl, quinoxalinyl, indoliziny, isoindolyl, benzothienyl, benzofuranyl, isobenzofuranyl and chromenyl groups.

48. (Withdrawn) The method of claim 1 wherein said compound has a general formula V-B



wherein

X<sub>5</sub>, Y<sub>3</sub>, and Z<sub>3</sub> are independently selected from the group consisting of N, O, S, C, and CO wherein at least one of X<sub>5</sub>, Y<sub>3</sub>, and Z<sub>3</sub> is a heteroatom;

Ar<sub>4</sub> and Ar<sub>5</sub> are independently selected from the group consisting heterocyclic and fused heterocyclic groups containing 1 to 4 heteroatoms selected from the group consisting of N, O, and S and an aromatic group selected from the group consisting of phenyl, benzyl, 1-naphthyl, 2-naphthyl, fluorenyl, anthrenyl, indenyl, phenanthrenyl, and benzonaphthenyl, wherein Ar<sub>4</sub> and Ar<sub>5</sub> are optionally substituted with one or more substituents selected from the group consisting of -F, -Cl, -Br, -I, -OR<sub>j</sub>, -SR<sub>j</sub>, -SOR<sub>j</sub>, -SO<sub>2</sub>R<sub>j</sub>, -SO<sub>2</sub>NR<sub>j</sub>R<sub>k</sub>, -OCOR<sub>j</sub>, -OCONR<sub>j</sub>R<sub>k</sub>, -NRCOR<sub>k</sub>, -NRCO<sub>2</sub>R<sub>k</sub>, -CN, -NO<sub>2</sub>, -CO<sub>2</sub>R<sub>j</sub>, -CONR<sub>j</sub>R<sub>k</sub>, -C(O)R<sub>j</sub>, -CH(OR<sub>j</sub>)R<sub>k</sub>, -CH<sub>2</sub>(OR<sub>j</sub>)-R<sub>j</sub>, and -A-(CH<sub>2</sub>)<sub>n</sub>-NR<sub>j</sub>R<sub>k</sub>; wherein R<sub>j</sub> and R<sub>k</sub> are selected independently from the group consisting of H, CF<sub>3</sub>, (C<sub>1-10</sub>)alkyl, cycloalkyl, alkyl-aryl, alkyl-heteroaryl, heterocycloalkyl, aryl, or R<sub>j</sub> and R<sub>k</sub> may combine to form a C<sub>1-5</sub> methylene chain, A is defined as CH<sub>2</sub>, O, NH, S, SO, SO<sub>2</sub>,

and n is 1, 2, 3, or 4,

or an enantiomer, diastereoisomer, N-oxide, crystalline form, hydrate, solvate, pharmacologically active metabolite, prodrug, or pharmaceutically acceptable salt thereof.

49. (Withdrawn) The method of claim 48 wherein said heterocyclic or fused heterocyclic group is selected from the group consisting of quinolyl, quinazolyl, quinoxalyl, 2-pyrimidyl, 4-pyrimidyl, 5-pyrimidyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, and pyrazyl.

50. (Withdrawn) A method of identifying a compound useful for treating neuromuscular dysfunction of the lower urinary tract in a mammal, comprising

- (a) determining the binding affinities of one or more test compound for an mGlu5 receptor and one or more of an mGlu1 receptor or Group II mGlu receptor;
- (b) identifying a test compound that
  - (1) binds to mGlu5 receptor with an affinity of at least  $10^{-6}$  M; and
  - (2) binds to mGlu5 receptor with an affinity at least 10-fold stronger than the affinity for mGlu1 receptor or Group II mGlu receptor.

51. (Withdrawn) The method of claim 50 further comprising  
individually measuring the binding affinity of said one or more test compounds for one or more Group III mGlu receptor and  
identifying a test compound that binds to mGlu5 receptor with an affinity at least 10-fold stronger than the affinity for a Group III mGlu receptor.

52. (Withdrawn) The method of claim 50 or 51 wherein step (b) comprises identifying a test compounds that

- (1) binds to mGlu5 receptor with an affinity of at least  $10^{-6}$  M; and
- (2) binds to mGlu5 receptor with an affinity at least 10-fold stronger than the affinity for each of mGlu1 receptor and Group II mGlu receptor.

53. (Withdrawn) The method of claim 50 or 51 wherein step (b) comprises identifying a test compounds that

- (1) binds to mGlu5 receptor with an affinity of at least  $10^{-6}$  M; and
- (2) binds to mGlu5 receptor with an affinity at least 100-fold stronger than the affinity for a mGlu1 receptor or Group II mGlu receptor.

54. (Withdrawn) The method of claim 53 wherein step (b) comprises identifying a test compounds that

- (1) binds to mGlu5 receptor with an affinity of at least  $10^{-6}$  M; and

